

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/99180/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Morris, Reginald ORCID: <https://orcid.org/0000-0002-8250-9315>, Eccles, Alicia, Ryan, Brooke and Kneebone, Ian I. 2017. The prevalence of anxiety in people with aphasia after stroke. *Aphasiology* 31 (12) , pp. 1410-1415.
10.1080/02687038.2017.1304633 file

Publishers page: <http://dx.doi.org/10.1080/02687038.2017.1304633>
< <http://dx.doi.org/10.1080/02687038.2017.1304633> >

Please note:


Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.





Prevalence of anxiety in people with aphasia after stroke

Reg Morris, Alicia Eccles, Brooke Ryan & Ian I. Kneebone

To cite this article: Reg Morris, Alicia Eccles, Brooke Ryan & Ian I. Kneebone (2017): Prevalence of anxiety in people with aphasia after stroke, Aphasiology, DOI: [10.1080/02687038.2017.1304633](https://doi.org/10.1080/02687038.2017.1304633)

To link to this article: <http://dx.doi.org/10.1080/02687038.2017.1304633>



© 2017 Informa UK Limited, trading as Taylor & Francis Group



Published online: 30 Mar 2017.



Submit your article to this journal [↗](#)



Article views: 77



View related articles [↗](#)



View Crossmark data [↗](#)

Prevalence of anxiety in people with aphasia after stroke

Reg Morris^a, Alicia Eccles^a, Brooke Ryan^b and Ian I. Kneebone^c

^aSchool of Psychology, University of Cardiff and Cardiff and Vale University Health Board, Cardiff, UK;

^bSchool of Health and Rehabilitation Sciences, The University of Queensland, Brisbane, Australia;

^cDiscipline of Clinical Psychology, Graduate School of Health, University of Technology Sydney, Ultimo, Australia

ABSTRACT

Background: Anxiety is common after stroke and is associated with poorer recovery. People with aphasia after stroke are typically excluded from studies of anxiety prevalence and so the number of those affected is unclear.

Aims: To make a preliminary estimate of the prevalence of significant anxiety in people with aphasia after stroke.

Methods & Procedures: Carers to community-dwelling people with aphasia after stroke, $N = 111$, completed the Behavioural Outcomes of Anxiety scale (BOA), a modified Hospital Anxiety and Depression Scale – Anxiety sub-scale (HADS-A), and a modified Generalised Anxiety Disorder-7 item (GAD-7) scale to determine the presence of significant anxiety in the person for whom they cared. Associates of anxiety in people with aphasia after stroke were also investigated.

Outcomes & Results: The BOA identified 49 people (44%) as having significant anxiety. Findings for the HADS-A = 46 (41%) were similar; however, for the GAD-7, the rate was substantially lower = 18 (16%). Anxiety after stroke had a modest but significant association with younger age (all measures) and with the Frenchay Aphasia Severity Test scores (BOA and HADS-A only).

Conclusions: The prevalence of anxiety in people with aphasia after stroke is high and likely higher than in those with stroke with no aphasia when measured using a validated tool for this population. Risk factors for anxiety appear to be severity of aphasia and younger age. This finding should be tempered by the fact that this is a preliminary study in a relatively small sample consisting of those attending stroke groups and the use of caregiver assessments may overestimate the prevalence of mood disorder.

ARTICLE HISTORY

Received 13 December 2016

Accepted 6 March 2017

KEYWORDS

Stroke; aphasia; anxiety; prevalence

Anxiety disorders is a class of conditions that includes generalised anxiety disorder, panic disorder, and phobias (American Psychiatric Association, 2013). Features common to these conditions are fear out of proportion, apprehension of danger or negative events, and day-to-day distress. Physical aspects can include tension, palpitations, dizziness, or trembling. These conditions are amongst the most common of psychiatric disorders. Post-traumatic stress disorder is amongst a class of conditions known as trauma- and stressor-related disorders and also commonly involves the experience of

significant anxiety symptoms (American Psychiatric Association, 2013). Stroke is typically a sudden onset life-threatening event and a leading cause of disability. It can impair physical, cognitive, and communication function (Lincoln, Kneebone, Macniven, & Morris, 2012). Unsurprisingly, stroke can be a precipitant to anxiety and post-traumatic stress symptoms. The point prevalence rate for significant anxiety after stroke is between 18% and 25% (Campbell Burton et al., 2013). While anxiety is challenging after stroke, of further concern is its impact on recovery. Anxiety after stroke appears to impact quality of life (Jeong et al., 2012; Tang, Lau, Mok, Ungvari, & Wong, 2013), social contact (Astrom, 1996), and functional ability (Astrom, 1996; D'Alisa, Baudo, Mauro, & Miscio, 2005). Aphasia affects between 30% and 34% of those with stroke (Flowers et al., 2016). Traditionally, significant anxiety is identified via questionnaire and/or clinical interview. On this account, those with aphasia are typically excluded from studies assessing anxiety prevalence (Campbell Burton et al., 2013). Preliminary research comparing small groups of those with and without aphasia after stroke provides some evidence to suggest anxiety is likely higher for those who have aphasia (Shehata, El Mistikawi, Al Sayed, & Hassan, 2015).

Recently, an instrument has been developed that has facilitated the assessment of anxiety in people with aphasia after stroke (Eccles, Morris, & Kneebone, 2016; Kneebone, Neffgen, & Pettyfer, 2012; Linley-Adams, Morris, & Kneebone, 2014). The Behavioural Outcomes of Anxiety (BOA) scale uses observer (typically formal or informal carer) ratings of behaviour to identify anxiety after stroke. The BOA allows anxiety prevalence to be assessed for the first time in this vulnerable group. It was the aim of the current study to specifically consider the prevalence of anxiety in community-dwelling people with aphasia after stroke.

Method

Participants were an opportunistic sample involved in the initial cross-sectional phase of a study to assess the psychometric properties of the BOA. A detailed description of the methodology is available elsewhere (Eccles et al., 2016). Participants with aphasia and their carers were recruited from stroke groups in Wales, United Kingdom. Inclusion criteria included having had a stroke between 2 months and 20 years previously and carer or self-reported aphasia. Communication was assessed using the Frenchay Aphasia Severity Test (FAST; Enderby, Wood, Wade, & Hewer, 1986). Any patient or carer <18 years old was excluded from participation.

Carers completed questionnaires that included the BOA, a carer-completed version of the Hospital Anxiety and Depression Scale-Anxiety sub-scale (HADS-A; Zigmond & Snaith, 1983) and a carer-completed version of the Generalised Anxiety Disorder-7 item (GAD-7; Spitzer, Kroenke, Williams, & Lowe, 2006). This was done either in their own home or privately at a stroke group venue.

Results

One hundred and eleven patient carer dyads were recruited. Approximately 69% of patients were male and 72% of carers were female. Patient mean age was 69.7 (SD = 10.7), for carers this was 64.7 (SD = 12.2). For 65 patients, it was their first stroke.

Table 1. Scores on anxiety measures and their correlates.

Measure	Mean (SD)	Range	Caseness	FAST	Patient age	Correlations carer age	Gender	Time since stroke (years)
BOA	15.12 (6.19)	1–28	49 (44%)	.20*	–.27*	–.20*	.03	–.07
HADS-A	6.90 (4.44)	0–20	46 (41%)	.20*	–.28*	–.15	.09	–.09
GAD-7	5.46 (5.38)	0–19	18 (16%)	.15	–.22*	–.10	.08	–.17

Notes. $n/N = 105-111$, BOA: Behavioural Outcomes of Anxiety, cut-off > 16, HADS-A: Hospital Anxiety and Depression Scale- Anxiety sub-scale, cut-off > 7, GAD-7: Generalised Anxiety Disorder-7 item, cut-off > 10, FAST: Frenchay Aphasia Severity Test. All correlations. Pearson's r , except Gender which uses η . * $p < .05$ level (1 tailed).

Mean FAST score was 19 (SD = 5.9). Aphasia severity category statistics (Dalemans, De Witte, Beurskens, Van Den Heuvel, & Wade, 2010) were as follows: 16 (14%) participants had severe aphasia (FAST score of 0–10), 43 (39%) participants had moderate aphasia (FAST score of 11–20), 46 (41%) had mild aphasia (FAST score 21–26), and 6 (5%) minimal aphasia (FAST score 27–30). Table 1 summarises the scores obtained for each of the anxiety measures and their correlates. A prevalence rate of up to 44% is indicated for anxiety in those with aphasia post-stroke. Anxiety was associated with younger age on all the measures of anxiety. Level of aphasia was modestly correlated with anxiety on two of the measures.

Discussion

For the first time, an estimate of the prevalence of significant anxiety in people with aphasia after stroke using an instrument specifically developed for this purpose is available. At 44%, the prevalence established by the BOA is high relative to anxiety in the non-aphasic post-stroke samples (Campbell Burton et al., 2013).

The finding of a lower prevalence of anxiety in this sample as measured by the GAD-7 is perplexing. However, unlike the HADS-A and the BOA, the GAD-7 has not been subject to evaluation in stroke. It may well be it is not an appropriate instrument to use in this population, or that a lower cut-off is more appropriate if it is to be used. A lower cut-off for those with stroke has been found to be required for other instruments assessing anxiety in stroke (e.g., Kneebone, Fife-Schaw, Lincoln, & Harder, 2015). Further, it should be noted that a score of >5 on the GAD-7 is considered indicative of “mild symptoms” of anxiety on this measure. At this cut-off, prevalence rates would be similar to those identified by the BOA and the HADS-A.

The modest correlation between the FAST scores and anxiety scores is consistent with findings that, in general, cognitive deficits are associated with anxiety after stroke (Menlove et al., 2015). The association of anxiety with younger age may indicate a risk factor to be considered in this population, though results from studies that have considered this in the whole stroke population are inconsistent, with some studies supporting it (Broomfield, Scoular, Welsh, Walters, & Evans, 2013; Castellanos-Pinedo et al., 2011) and others not (Lincoln et al., 2013; Tang et al., 2012).

Some limitations to the current study should be acknowledged. First, this was a relatively small sample on which to estimate prevalence and also one that might not be fully representative of those with aphasia after stroke. As all participants were attenders at a stroke

group, they may be those who have sought out a group on account of anxiety, resulting in an overestimate of anxiety prevalence. In contrast, the study findings might underestimate anxiety on this basis of accessing participants from a stroke groups; those with anxiety might be less likely to attend a stroke group given both aphasia and anxiety after stroke is associated with social isolation (Astrom, 1996; Hilari & Northcott, 2016). Second, there is evidence from work considering depression in people after stroke that suggests carers might overestimate the presence of symptoms (Berg, Lönnqvist, Palomäki, & Kaste, 2009). There is some support for this contention in the work that took place in the development of the BOA – carers estimated anxiety as higher than did the stroke patient themselves (Linley-Adams et al., 2014). Finally, while screens such as the BOA detect anxiety symptoms, in themselves, they are not diagnostic of a particular disorder, for example, generalised anxiety disorder or PTSD. This should be considered if the interest is in estimating disorder-specific prevalence.

Robust population-based studies are required to support the currently suggested high rate of anxiety symptoms that appears evident in people with aphasia after stroke. Such studies should control for a possible contamination effect of anxiety in carers that complete the ratings and also consider to what extent a particular screening instrument might indicate the presence or absence of a particular psychiatric disorder in which anxiety symptoms may be prominent. Comprehensive specialist speech and language assessment of those with aphasia and anxiety after stroke might also improve an understanding of the relationship between communication disorder and anxiety in this population.

As anxiety is distressing, challenging and appears to affect function, it is a substantial problem worthy of treatment attention. This might be medication and/or psychological therapy. One psychological therapy, behavioural activation, has already proven useful in treating people with aphasia who have depression (Thomas, Walker, Macniven, Haworth, & Lincoln, 2013) and there is preliminary evidence relaxation training might be useful to treat anxiety in this population (Eccles et al., 2016). A modified cognitive behaviour therapy that takes into account communication and cognitive disabilities has also been proposed for people with stroke (Kneebone, 2016), and there is casework support for this with respect to anxiety management (Kneebone & Jeffries, 2013).

Acknowledgements

The authors gratefully acknowledge the support of The Stroke Association, UK, and the co-operation of the patients and carers who participated. Anastasia Hronis provided editorial assistance.

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

This research was made possible by funding from the Workforce and Education Development Service of the Welsh Government.

References

- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Washington, DC: American Psychiatric Association.
- Astrom, M. (1996). Generalized anxiety disorder in stroke patients: A 3-year longitudinal study. *Stroke*, 27, 270–275. doi:[10.1161/01.STR.27.2.270](https://doi.org/10.1161/01.STR.27.2.270)
- Berg, A., Lönnqvist, J., Palomäki, H., & Kaste, M. (2009). Assessment of depression after stroke: A comparison of different screening instruments. *Stroke*, 40, 523–529. doi:[10.1161/STROKEAHA.108.527705](https://doi.org/10.1161/STROKEAHA.108.527705)
- Broomfield, N., Scoular, A., Welsh, P., Walters, M., & Evans, J. J. (2013). Post-stroke anxiety is prevalent at the population level, especially among socially deprived and younger age community stroke survivors. *International Journal of Stroke*. doi:[10.1111/ijvs.12109](https://doi.org/10.1111/ijvs.12109)
- Campbell Burton, C. A., Murray, J., Holmes, J., Astin, F., Greenwood, D., & Knapp, P. (2013). Frequency of anxiety after stroke: A systematic review and meta-analysis of observational studies. *International Journal of Stroke*, 8, 545–559. doi:[10.1111/j.1747-4949.2012.00906.x](https://doi.org/10.1111/j.1747-4949.2012.00906.x)
- Castellanos-Pinedo, F., Hernández-Pérez, J. M., Zurdo, M., Rodríguez-Fúnez, B., Hernández-Bayo, J. M., García-Fernández, C., ... Castro-Posada, J. A. (2011). Influence of premorbid psychopathology and lesion location on affective and behavioral disorders after ischemic stroke. *Journal of Neuropsychiatry and Clinical Neurosciences*, 23, 340–347. doi:[10.1176/jnp.23.3.jnp340](https://doi.org/10.1176/jnp.23.3.jnp340)
- D'Alisa, S., Baudo, S., Mauro, A., & Miscio, G. (2005). How does stroke restrict participation in long-term post-stroke survivors? *Acta Neurologica Scandinavica*, 112, 157–162. doi:[10.1111/j.1600-0404.2005.00466.x](https://doi.org/10.1111/j.1600-0404.2005.00466.x)
- Dalemans, R. J., De Witte, L. P., Beurskens, A. J., Van Den Heuvel, W. J., & Wade, D. T. (2010). Psychometric properties of the community integration questionnaire adjusted for people with aphasia. *Archives of Physical Medicine and Rehabilitation*, 91, 395–399. doi:[10.1016/j.apmr.2009.10.021](https://doi.org/10.1016/j.apmr.2009.10.021)
- Eccles, A., Morris, R., & Kneebone, I. I. (2016). Psychometric properties of the Behavioural Outcomes of Anxiety questionnaire in stroke patients with aphasia. *Clinical Rehabilitation*. doi:[10.1177/0269215516644311](https://doi.org/10.1177/0269215516644311)
- Enderby, P. M., Wood, V. A., Wade, D. T., & Hewer, R. L. (1986). The Frenchay Aphasia Screening Test: A short, simple test for aphasia appropriate for non-specialists. *International Rehabilitation Medicine*, 8, 166–170. doi:[10.3109/03790798709166209](https://doi.org/10.3109/03790798709166209)
- Flowers, H. L., Skoretz, S. A., Silver, F. L., Rochon, E., Fang, J., Flamand-Roze, C., & Martino, R. (2016). Poststroke aphasia frequency, recovery and outcomes: A systematic review and meta-analysis. *Archives of Physical Medicine and Rehabilitation*, 97, 2188–2201.e8. doi:[10.1016/j.apmr.2016.03.006](https://doi.org/10.1016/j.apmr.2016.03.006)
- Hilari, K., & Northcott, S. (2016). “Struggling to stay connected”: Comparing the social relationships of healthy older people and people with stroke and aphasia. *Aphasiology*, 1–14. doi:[10.1080/02687038.2016.1218436](https://doi.org/10.1080/02687038.2016.1218436)
- Jeong, B. O., Kang, H. J., Bae, K. Y., Kim, S. W., Kim, J. M., Shin, I. S., ... Yoon, J. S. (2012). Determinants of quality of life in the acute stage following stroke. *Psychiatry Investigation*, 9, 127–133. doi:[10.4306/pi.2012.9.2.127](https://doi.org/10.4306/pi.2012.9.2.127)
- Kneebone, I. I. (2016). A framework to support cognitive behavior therapy for emotional disorder after stroke. *Cognitive and Behavioral Practice*, 23, 99–109. doi:[10.1016/j.cbpra.2015.02.001](https://doi.org/10.1016/j.cbpra.2015.02.001)
- Kneebone, I. I., Fife-Schaw, C., Lincoln, N. B., & Harder, H. (2015). A study of the validity and the reliability of the Geriatric Anxiety Inventory in screening for anxiety after stroke in older inpatients. *Clinical Rehabilitation*. doi:[10.1177/0269215515619661](https://doi.org/10.1177/0269215515619661)
- Kneebone, I. I., & Jeffries, F. W. (2013). Treating anxiety after stroke using cognitive behaviour therapy: Two cases. *Neuropsychological Rehabilitation*, 23, 798–810. doi:[10.1080/09602011.2013.820135](https://doi.org/10.1080/09602011.2013.820135)
- Kneebone, I. I., Neffgen, L., & Pettyfer, S. (2012). Screening for depression and anxiety after stroke: Developing protocols for use in the community. *Disability and Rehabilitation*, 34, 1114–1120. doi:[10.3109/09638288.2011.636137](https://doi.org/10.3109/09638288.2011.636137)

- Lincoln, N., Kneebone, I. I., Macniven, J., & Morris, R. (2012). *Psychological management of stroke*. Chichester, UK: Wiley.
- Lincoln, N. B., Brinkmann, N., Cunningham, S., Dejaeger, E., De Weerd, W., Jenni, W., ... De Wit, L. (2013). Anxiety and depression after stroke: A 5 year follow up. *Disability and Rehabilitation*, 35, 140–145. doi:[10.3109/09638288.2012.691939](https://doi.org/10.3109/09638288.2012.691939)
- Linley-Adams, B., Morris, R., & Kneebone, I. (2014). The Behavioural Outcomes of Anxiety scale (BOA): A preliminary evaluation in stroke survivors. *British Journal of Clinical Psychology*, 53, 451–467. doi:[10.1111/bjc.12056](https://doi.org/10.1111/bjc.12056)
- Menlove, L., Crayton, E., Kneebone, I., Allen-Crooks, R., Otto, E., & Harder, H. (2015). Predictors of anxiety after stroke: A systematic review of observational studies. *Journal of Stroke and Cerebrovascular Diseases*, 24, 1107–1117. doi:[10.1016/j.jstrokecerebrovasdis.2014.12.036](https://doi.org/10.1016/j.jstrokecerebrovasdis.2014.12.036)
- Shehata, G. A., El Mistikawi, T., Al Sayed, K. R., & Hassan, H. S. (2015). The effect of aphasia upon personality traits, depression and anxiety among stroke patients. *Journal of Affective Disorders*, 172, 312–314. doi:[10.1016/j.jad.2014.10.027](https://doi.org/10.1016/j.jad.2014.10.027)
- Spitzer, R. L., Kroenke, K., Williams, J. B., & Lowe, B. (2006). A brief measure for assessing generalised anxiety disorder: The Generalised Anxiety Disorder-7. *Archives of Internal Medicine*, 166, 1092–1097. doi:[10.1001/archinte.166.10.1092](https://doi.org/10.1001/archinte.166.10.1092)
- Tang, W. K., Chen, Y., Lu, J., Liang, H., Chu, W. C. W., Mok, V. C. T., ... Wong, K. S. (2012). Frontal infarcts and anxiety in stroke. *Stroke*, 43, 1426–1428. doi:[10.1161/STROKEAHA.111.640482](https://doi.org/10.1161/STROKEAHA.111.640482)
- Tang, W. K., Lau, C. G., Mok, V., Ungvari, G. S., & Wong, K. S. (2013). Impact of anxiety on health-related quality of life after stroke: A cross-sectional study. *Archives of Physical Medicine and Rehabilitation*, 94, 2535–2541. doi:[10.1016/j.apmr.2013.07.012](https://doi.org/10.1016/j.apmr.2013.07.012)
- Thomas, S. A., Walker, M. F., Macniven, J. A., Haworth, H., & Lincoln, N. B. (2013). Communication and Low Mood (CALM): A randomized controlled trial of behavioural therapy for stroke patients with aphasia. *Clinical Rehabilitation*, 27, 398–408. doi:[10.1177/0269215512462227](https://doi.org/10.1177/0269215512462227)
- Zigmond, A. S., & Snaith, R. P. (1983). Hospital Anxiety and Depression Scale. *Acta Psychiatrica Scandinavica*, 67, 361–370. doi:[10.1111/j.1600-0447.1983.tb09716.x](https://doi.org/10.1111/j.1600-0447.1983.tb09716.x)